

Abelian representation for nonabelian Wilson loops and the Non - Abelian Stokes theorem on the lattice.

M.A.Zubkov*

ITEP, B.Chermushkinskaya 25, Moscow, 117259, Russia

(Dated: July, 12, 2003)

We derive the Abelian - like expression for the lattice $SU(N)$ Wilson loop in arbitrary irreducible representation. The continuum Abelian representation of the $SU(N)$ Wilson loop (for the loop without selfintersections) that has been obtained by Diakonov and Petrov appears to be a continuum limit of this expression. We also obtain the lattice variant of a non - Abelian Stokes theorem and present the explicit expression for the matrix $\mathcal{H}$ used in the Diakonov - Petrov approach.

PACS numbers: 11.15.Ha, 12.38.Lg, 12.38.Gc

I. INTRODUCTION

In order to visualize multidimensional space the projection onto the lower dimensional spaces can be used. The analogous idea is used in the non - Abelian gauge theory, where the dynamics is described in terms of certain Abelian variables. This method is called the Abelian Projection [1].

One of the most important quantities we encounter in gauge theory is the Wilson loop

$$W_q[\mathcal{C}] = \chi_q(\text{Pexp}(i \int_{\mathcal{C}} A_{\mu} dx_{\mu})) \quad (1)$$

Here A is the gauge field (that belongs to the Lie algebra of the correspondent gauge group), $\mathcal{C}$ is the closed contour, P denotes the usual path ordering, and χ_q is the character considered in the representation q of the gauge group. For the application of the Abelian projection method it is important to have an Abelian - like representation for the Wilson loop. The representation of such kind was derived more than ten years ago by D. Diakonov and V. Petrov [3]. For the case of the irreducible representation q of $SU(N)$ it has the form:

$$W_q[\mathcal{C}] = \int D\mu_{\mathcal{C},q}(g) \exp(i \int_{\mathcal{C}} \text{Tr} A_{\mu}^g \mathcal{H}^q dx_{\mu}), \quad (2)$$

where g is the $SU(N)$ gauge transformation and $A_{\mu}^g = g A_{\mu} g^+ - i g \partial_{\mu} g^+$. Here $\mathcal{H}^q = \sum_{i=1, \dots, N-1} m_i H_i$, where each H_i denotes the basis element of the Cartan subalgebra of $su(N)$ and (m_i) is the highest weight of the representation. The matrices H_i are normalized in such a way that $\text{Tr} H_i H_j = \delta_{ij}$. The gauge transformations are defined on the contour $\mathcal{C}$. The measure $\mu_{\mathcal{C},q}(g)$ on the space of gauge transformations is constructed using the invariant measure on the $SU(N)$ group. This construction will be discussed in detail later.

Both in the monopole and in the P - vortex confinement pictures the confining forces are induced by the magnetic flux contained inside the contour that is correspondent to the worldline of the charged particle [2]. In the Abelian theory a Stokes theorem is used for counting the magnetic flux. For counting the "magnetic flux" contained inside the contour correspondent to the Wilson loop in a non - Abelian theory D. Diakonov and V. Petrov suggested to use representation (2) in the following form:

$$W_q[\mathcal{C}] = \int D\mu_{\mathcal{C},q}(g) \exp(i \int_{\mathcal{M}(\mathcal{C})} \text{Tr} \partial_{[\mu} A_{\nu]}^g \mathcal{H}^q dx_{\mu} \wedge dx_{\nu}), \quad (3)$$

where g is defined on the surface $\mathcal{M}(\mathcal{C})$ that spans the contour $\mathcal{C}$. The integral of a two - form over this surface is denoted by $\int_{\mathcal{M}(\mathcal{C})} \dots dx_{\mu} \wedge dx_{\nu}$. The expression (3) is known as the non - Abelian Stokes theorem.

In the present work we consider a lattice variant of the Abelian representation for the non - Abelian Wilson loop and the non - Abelian Stokes theorem on the lattice. The representation (2) was originally derived in a continuum theory. In the later publications (see, for example, [4, 5]) the approximated lattice variant of (2) was derived.

*Electronic address: zubkov@itep.ru

It appears that one of the most fruitful methods of the investigation of the quantum field theory is the lattice approach. This approach gives the exact meaning to any field correlator. It is possible both to define and to calculate it using the lattice regularization. Without using a lattice numerical investigation this could become possible only for the Integrable Models.

Thus it seems important to derive the lattice variant of (2), which is valid within the lattice $SU(N)$ theory *precisely* (i.e. not only in its naive continuum limit). In this paper such a representation is derived. If we assume that the gauge field is smooth enough, then the continuum limit of the derived representation coincides with (2). The non-Abelian Stokes theorem on the lattice commonly follows from the lattice Abelian representation of the Wilson loop.

II. IRREDUCIBLE REPRESENTATIONS OF THE $SU(N)$ GROUP

In this section we briefly summarize the basic notations of group representation theory that will be used in the next sections.

1. The space $\mathcal{V}$ of any irreducible representation of $SU(N)$ consists of the tensors $\Psi_{i_1 i_2 \dots i_r}$. The symmetry of Ψ is defined by the set of integer numbers q_i ($i = 1, \dots, N-1$) ($\sum_i q_i = r$, $q_i \geq 0$), which is called the signature of the representation. The elements of $\mathcal{V}$ are the tensors

$$\Psi_{i_1^1 \dots i_{q_1}^1 \dots i_1^{c_1} \dots i_{q_{c_1}}^{c_1}} \quad (4)$$

For any $k \in \{1, \dots, q_1\}$ the number of indices i_l^k is denoted by c_k . Let us arrange these indices as follows:

$$\begin{array}{cccc} i_1^1 & \dots & i_i^1 & \dots & i_{q_1}^1 \\ i_1^2 & \dots & i_{q_2}^2 & & \\ \dots & & & & \\ i_1^{c_1} & \dots & & & \end{array} \quad (5)$$

Therefore q_i is the length of the i -th line while the height of the l -th column is c_l .

2. Let us consider the arbitrary tensor $\Psi_{i_1^1 \dots i_{q_1}^1 \dots i_1^{c_1} \dots i_{q_{c_1}}^{c_1}}$. The tensor with the required symmetry can be obtained by means of the following transformation:

$$\begin{aligned} \Psi_{i_1^1 \dots i_{q_1}^1 \dots i_1^{c_1} \dots i_{q_{c_1}}^{c_1}} &\rightarrow \mathcal{S}^q \Psi_{i_1^1 \dots i_{q_1}^1 \dots i_1^{c_1} \dots i_{q_{c_1}}^{c_1}} \\ &= \sum_S (-)^{P(S)} \Psi_{S[i_1^1 \dots i_{q_1}^1 \dots i_1^{c_1} \dots i_{q_{c_1}}^{c_1}]}, \end{aligned} \quad (6)$$

where the sum is over the permutations S of the indices. Each permutation that enters the sum consists of the ordered product $\mathcal{Q} \times \mathcal{P}$ of some permutation $\mathcal{P}$ inside the lines $(i_1^k \dots i_{q_k}^k)$ of (5) and some permutation $\mathcal{Q}$ inside the columns $(i_l^1 \dots i_l^{c_l})$. $P(S) = 0(1)$ if Q is even (odd).

3. Let U_{ij} be the $N \times N$ matrix that represents a certain element of $SU(N)$. Then the action of this element on $\mathcal{V}$ is given by:

$$\begin{aligned} \Psi_{i_1 i_2 \dots i_r} &\rightarrow \mathcal{D}_q(U)_{i_1 i_2 \dots i_r}^{j_1 j_2 \dots j_r} \Psi_{j_1 j_2 \dots j_r} \\ &= U_{i_1 j_1} U_{i_2 j_2} \dots U_{i_r j_r} \Psi_{j_1 j_2 \dots j_r} \end{aligned} \quad (7)$$

We shall also write:

$$|\Psi\rangle \rightarrow \mathcal{D}_q(U)|\Psi\rangle \quad (8)$$

4. Let us represent an infinitesimal element of $SU(N)$ as $U = \exp(iAs)$. Here s is an infinitely small parameter and A is a matrix that represents a certain element of the Lie algebra $su(N)$. Then (7) defines the representation of $su(N)$. We denote the correspondent matrix by $\mathcal{D}_q(A)$:

$$\mathcal{D}_q(U) = \exp(i\mathcal{D}_q(A)s) \quad (9)$$

5. Using the Cartan basis any $A \in su(N)$ can be represented as

$$\begin{aligned} A &= \sum_{i=1, \dots, N-1} H^i \text{Tr} A H^i + \\ &\sum_{i < j; i, j=1, \dots, N-1} (E^{ij} (\text{Tr} A E^{ij})^* + (E^{ij})^+ \text{Tr} A E^{ij}) \end{aligned} \quad (10)$$

The Cartan elements of the basis are

$$H^i = \frac{1}{\sqrt{i(i+1)}} \text{diag}(1, \dots, 1, -i, 0, \dots) \quad (11)$$

The shift operators are:

$$(E^{ij})_{ab} = \delta_{ai}\delta_{bj} \quad (12)$$

6. The normalized vector $|\Lambda^a\rangle$ ($\langle\Lambda^a|\Lambda^a\rangle = 1$), which is an eigenvector for any operator $\mathcal{D}_q(H^i)$ is called the weight vector. Here Λ^a ($a = 1, \dots, N-1$) are the correspondent eigenvalues:

$$\mathcal{D}_q(H^i)|\Lambda^a\rangle = \Lambda^i|\Lambda^a\rangle \quad (13)$$

Shift operators act on the weight vectors as follows:

$$\begin{aligned} \mathcal{D}_q(E^{ij})|\Lambda^a\rangle &\sim |\Lambda^a + (\alpha^{ij})^a\rangle \\ \mathcal{D}_q(E^{ij})^+|\Lambda^a\rangle &\sim |\Lambda^a - (\alpha^{ij})^a\rangle \end{aligned} \quad (14)$$

Here $|a\rangle \sim |b\rangle$ means that $|a\rangle = C|b\rangle$ for some constant C . The numbers α^{ij} are called the roots of $su(N)$. They enter the following commutation relations:

$$\begin{aligned} [H^a, E^{ij}] &= (\alpha^{ij})^a E^{ij} \\ [H^a, (E^{ij})^+] &= -(\alpha^{ij})^a (E^{ij})^+ \end{aligned} \quad (15)$$

7. The weight vector that is annihilated by each of the operators E^{ij} is called the eldest vector. The correspondent set of numbers Λ^a is called the highest weight and is denoted by m^a .

We denote the eldest vector by $|0\rangle$. Up to the normalization factor it corresponds to the tensor

$$\Psi_{i_1^1 \dots i_{q_1}^1 \dots i_{q_{c_1}}^1 \dots i_{q_{c_1}}^{c_1}}^0 \sim \mathcal{S}^q \delta_{i_1^1}^1 \dots \delta_{i_{q_1}^1}^1 \dots \delta_{i_{q_{c_1}}^1}^{c_1} \dots \delta_{i_{q_{c_1}}^{c_1}}^{c_1} \quad (16)$$

8. The highest weight of the representation q is given by:

$$m_i = \frac{1}{\sqrt{i(i+1)}} (q_1 + q_2 + \dots + q_{i-1} - iq_i) \quad (17)$$

9. The coherent state system correspondent to the representation q is the following subset of $\mathcal{V}$:

$$\{|g\rangle : |g\rangle = e^{i\phi(g)} \mathcal{D}_q(g)|0\rangle \ (g \in SU(N))\}, \quad (18)$$

where $\phi(g)$ is a real - valued function on $SU(N)$.

10. The coherent state system possesses the following completeness relation:

$$\int dg |g\rangle \langle g| = \frac{1}{D(q)} \quad (19)$$

Here $D(q)$ is the dimension of the representation. The measure on $SU(N)$ is normalized in such a way that $\int dg = 1$.

III. THE ABELIAN REPRESENTATION FOR THE NON - ABELIAN WILSON LOOP ON THE LATTICE.

In this section we consider the lattice $SU(N)$ Wilson loop. As an auxiliary instrument we use the lattice $SU(N)$ gauge model coupled to the scalar field Φ that belongs to the irreducible representation q of $SU(N)$. In addition this scalar field is considered in the presence of the external $U(1)$ gauge field $\theta \in]-\pi, \pi]$ (that lives on the links of the lattice). The partition function has the form:

$$Z[\theta] = \langle \int D\Phi \exp(\sum_{xy} \Phi_x^+ \mathcal{D}_q(U_{xy}) e^{i\theta} \Phi_y - V(\Phi)) \rangle \quad (20)$$

Here $U \in SU(N)$ and the average is calculated in a certain $SU(N)$ gauge theory. We assume that the potential V is infinitely deep and that it has the minimum at $\Phi = |0\rangle$ with $V(|0\rangle) = 0$, where $|0\rangle$ is the eldest vector of the representation. We also assume that V is invariant under the transformation $\Phi \rightarrow e^{i\alpha} \mathcal{D}_q(g)\Phi$, where α is arbitrary real-valued field.

The idea is that there exist two different ways to rewrite the partition function as a sum over the worldlines of the scalar. The first one demonstrates that the partition function of the model generates the non - Abelian Wilson loop average (in the representation q). The second one demonstrates that the same functional also generates the Abelian - like expression for the Wilson loop. Finally due to the arbitrariness of the $SU(N)$ measure we obtain the equivalence of the Wilson loop itself and its Abelian - like representation.

A. $Z[\theta]$ generates the Wilson loops

The mentioned properties of the potential make it possible to reduce the integration over Φ to the integration over $g \in SU(N)$ and over a real - valued scalar field α . Both fields are defined on the sites of the lattice. Then we can rewrite (20) as a sum over the worldlines of the scalar as follows:

$$\begin{aligned} Z[\theta] &= \langle \int D\alpha Dg \exp(\sum_{xy} \langle 0 | \mathcal{D}_q(g)_x^+ \mathcal{D}_q(U_{xy}) \\ &\quad e^{i\theta_{xy} + i\alpha_y - i\alpha_x} \mathcal{D}_q(g)_y | 0 \rangle) \rangle \\ &= \langle \int Dg \sum_{\mathcal{C}} (2\pi)^{n(j(\mathcal{C}))} e^{i(j(\mathcal{C}), \theta)} \\ &\quad \text{Tr} \Pi_{\{xy\} \in \mathcal{C}} \frac{1}{|j(\mathcal{C})_{xy}|!} \langle g_x | \mathcal{D}_q(U_{xy}) | g_y \rangle^{|j(\mathcal{C})_{xy}|} \rangle \end{aligned} \quad (21)$$

Here we used the calculus of differential forms on the lattice (for the definitions see, for example, [8]). The sum is over the closed lattice paths $\mathcal{C}$. The closed lattice path is defined as a map $\mathcal{C} : Z_N = \{0, 1, \dots, N-1\} \rightarrow X$ (where X is a set of the lattice points and N is an integer number) such that for each $k \in Z_N$ the points $\mathcal{C}(k)$ and $\mathcal{C}([k+1] \bmod N)$ are connected by a link. The product over all such links is denoted by $\Pi_{\{xy\} \in \mathcal{C}}$, where each link is counted once and it is implied that $x = \mathcal{C}(k)$ and $y = \mathcal{C}([k+1] \bmod N)$ for some k . $j(\mathcal{C})$ is an integer - valued link field that corresponds to the path $\mathcal{C}$. $|j_{xy}|$ counts the number of times the given link $\{xy\}$ is encountered when we are moving along the path. The sign of j_{xy} reflects the orientation of the path $\mathcal{C}$. So the case $|j| > 1$ corresponds to the overlapping of the different pieces of $\mathcal{C}$. It is worth mentioning that j does not contain the complete information on the orientation of $\mathcal{C}$. Therefore different paths may correspond to the same 1 - form $j(\mathcal{C})$. This may happen if $\mathcal{C}$ has the selfintersections. However, in (21) the sum is over $\mathcal{C}$ that have *different* $j(\mathcal{C})$. This means that the sum is indeed over different $j(\mathcal{C})$ and each term of the sum depends only upon $j(\mathcal{C})$ (not on $\mathcal{C}$ itself). $n(j)$ is the number of points, which enter the path represented by j . Also we assume the following notation:

$$(j, \theta) = \sum_{xy} j_{xy} \theta_{xy}, \quad (22)$$

where the sum is over the links of the lattice.

We denote $\langle i | \mathcal{D}_q(g) | 0 \rangle = \mathcal{D}_q(g)_{i0}$; $\langle 0 | \mathcal{D}_q(g)^+ | i \rangle = \mathcal{D}_q(g)_{0i}^+$, where $|i\rangle$ is the basis vector of the space $\mathcal{V}$ of the representation q . The integration over Dg can be performed using the following expressions:

$$\begin{aligned} \int dg \mathcal{D}_q(g)_{i0}^+ \mathcal{D}_q(g)_{0j} &= \frac{1}{M_1(q)} \delta_{ij} \\ \int dg \mathcal{D}_q(g)_{i_1 0}^+ \mathcal{D}_q(g)_{0 j_1} \mathcal{D}_q(g)_{i_2 0}^+ \mathcal{D}_q(g)_{0 j_2} &= \\ \frac{1}{M_2(q)} (\delta_{i_1 j_1} \delta_{i_2 j_2} + \delta_{i_1 j_2} \delta_{i_2 j_1}) & \\ \dots & \end{aligned} \quad (23)$$

Here the measure dg is normalized in such a way that $\int dg = 1$. The first identity reflects the completeness of a coherent state system and implies that $M_1(q) = D(q)$, where $D(q)$ is the dimension of the representation. Other equations follow from the gauge invariance. The correspondent coefficients $M_i(q)$ depend on the representation and

can be calculated explicitly using the expressions for the basis vectors of $\mathcal{V}$ and the technique of a strong coupling expansion (see, for example, [9]).

We obtain:

$$Z[\theta] = \langle \sum_{\mathcal{C}} C(j(\mathcal{C})) e^{i(j(\mathcal{C}), \theta)} W_q[\mathcal{C}] \rangle \quad (24)$$

where the sum is over *all different* $\mathcal{C}$. $W_q[\mathcal{C}]$ is the Wilson loop (in the representation q) correspondent to the path $\mathcal{C} : Z_N \rightarrow X$:

$$W_q[\mathcal{C}] = \text{Tr} \mathcal{D}_q(U_{\mathcal{C}(0)\mathcal{C}(1)}) \mathcal{D}_q(U_{\mathcal{C}(1)\mathcal{C}(2)}) \dots \mathcal{D}_q(U_{\mathcal{C}(N-1)\mathcal{C}(0)}) \quad (25)$$

We also denote

$$C(j) = \frac{(2\pi)^{n(j(\mathcal{C}))}}{(M_1(q))^{n_1(j)} (M_2(q))^{n_2(j)} \dots} \quad (26)$$

Here n_1 is the number of points of the path $\mathcal{C}$, where there are no selfintersections, n_2 is the number of points where $\mathcal{C}$ intersects itself once (i.e. where there are two different $k_a \in Z_N$, ($a = 1, 2$), such that $\mathcal{C}(k_1) = \mathcal{C}(k_2)$), n_2 is the number of points where $\mathcal{C}$ intersects itself twice etc ($n = \sum_i n_i$).

As we have already mentioned, j does not contain the complete information on the orientation of $\mathcal{C}$. Hence we extract from (24) the sum of the Wilson loops, which have the same $j(\mathcal{C})$. The expression for the vacuum average of $W_q^l = \sum_{j[\mathcal{C}] = l} W_q[\mathcal{C}]$ can be obtained using the Fourier transformation:

$$\begin{aligned} \langle W_q^l \rangle &= \sum_{j[\mathcal{C}] = l} \langle W_q[\mathcal{C}] \rangle \\ &= \frac{1}{C[l]} \int D\theta \exp(-i(l, \theta)) Z[\theta] \end{aligned} \quad (27)$$

B. $Z[\theta]$ generates the Abelian - like expressions for the Wilson loop

Let us fix the gauge, where $\Phi_x = e^{i\alpha_x} |0\rangle$ (α is a real valued field). We get:

$$\begin{aligned} Z[\theta] &= \langle \int D\alpha \exp(\sum_{xy} \langle 0 | \mathcal{D}_q(U_{xy}) | 0 \rangle e^{i(\theta_{xy} + \alpha_x - \alpha_y)}) \rangle \\ &= \langle \sum_{\mathcal{C}} \bar{C}(j(\mathcal{C})) \exp(i(j(\mathcal{C}), \theta)) \rangle \\ &= \Pi_{\{xy\} \in \mathcal{C}} \frac{1}{|j(\mathcal{C})_{xy}|!} \langle 0 | \mathcal{D}_q(U_{xy}) | 0 \rangle^{|j(\mathcal{C})_{xy}|} \end{aligned} \quad (28)$$

Here the sum is over $\mathcal{C}$ that have different $j(\mathcal{C})$ and

$$\bar{C}(j) = (2\pi)^{n(j)} \quad (29)$$

C. Equivalence of the Wilson loop and its Abelian representation

Recall that (24) and (28) are the different representations of the same quantity. Hence we obtain:

$$\begin{aligned} \langle W_q^l \rangle &= \frac{1}{C(l)} \int D\theta \exp(-i(l, \theta)) Z[\theta] \\ &= \sum_{j[\mathcal{C}] = l} \frac{\bar{C}(l)}{C(l)} \langle \Pi_{\{xy\} \in \mathcal{C}} \frac{1}{|j(\mathcal{C})_{xy}|!} \langle 0 | \mathcal{D}_q(U_{xy}) | 0 \rangle^{|j(\mathcal{C})_{xy}|} \rangle \\ &= \sum_{j[\mathcal{C}] = l} \tilde{C}(l) \langle \int Dg \Pi_{\{xy\} \in \mathcal{C}} \langle 0 | \mathcal{D}_q(U_{xy}^g) | 0 \rangle^{|j(\mathcal{C})_{xy}|} \rangle, \end{aligned} \quad (30)$$

where we used the gauge invariance of the operation $\langle \dots \rangle$ and introduced a new combinatorial factor

$$\tilde{C}(l) = (M_1(q)^{n_1(l)} M_2(q)^{n_2(l)} \dots) \prod_{\text{links}} \frac{1}{|l_{\text{link}}|!}, \quad (31)$$

where the sum is over all links of the lattice. The gauge transformation of the link variable U_{xy} is denoted by $U_{xy}^g = g U_{xy} g^+$. The measure Dg is defined in such a way that $\int Dg = 1$.

Due to the arbitrariness of the averaging $\langle \dots \rangle$ over the gauge field U we have:

$$W_q^l = \sum_{j[\mathcal{C}] = l} \tilde{C}[l] \int Dg \Pi_{\{xy\} \in \mathcal{C}} \langle 0 | \mathcal{D}_q(U_{xy}^g) | 0 \rangle^{|l_{xy}|} \quad (32)$$

Thus the sum of $SU(N)$ Wilson loops that are correspondent to the same 1 - form l can be represented through the Abelian - like expressions $\Pi_{\{xy\} \in \mathcal{C}} \langle 0 | \mathcal{D}_q(U_{xy}^g) | 0 \rangle^{|l_{xy}|}$. For the Wilson loops without selfintersections we have:

$$W^{l[\mathcal{C}]} = W[\mathcal{C}] \quad (33)$$

Then $\tilde{C}[l] = D(q)^{n(l)} = D(q)^{|\mathcal{C}|}$, where $|\mathcal{C}|$ is the length of the loop. In this case (32) acquires the form:

$$W_q[\mathcal{C}] = D(q)^{|\mathcal{C}|} \int Dg \Pi_{xy \in \mathcal{C}} \langle 0 | \mathcal{D}_q(U_{xy}^g) | 0 \rangle, \quad (34)$$

The expression (34) can be considered as the main result of this paper. In the next section we shall obtain the explicit expression for the matrix elements entering (34) and discuss symmetry properties of (34).

IV. PROPERTIES OF THE DERIVED REPRESENTATION.

In this section we derive the expression for $\langle 0 | \mathcal{D}_q(U_{xy}^g) | 0 \rangle$ entering (34). Also we introduce some new notations in order to demonstrate the Abelian nature of (34).

We use the explicit expressions for the eldest vector and for the matrix $\mathcal{D}_q(U)$ given in the section 2. We can represent $\langle 0 | \mathcal{D}_q(U) | 0 \rangle$ as

$$\langle 0 | \mathcal{D}_q(U) | 0 \rangle = \mathcal{U}(U), \quad (35)$$

where

$$\begin{aligned} \mathcal{U}(U) = \frac{1}{K} \sum_{S, \bar{S}} (-)^{P(S) + P(\bar{S})} U_{\bar{S}_1^1 S_1^1} \dots U_{\bar{S}_{q_1}^1 S_{q_1}^1} \\ \dots U_{\bar{S}_1^{c_1} S_1^{c_1}} \dots U_{\bar{S}_{q_{c_1}}^{c_1} S_{q_{c_1}}^{c_1}} \end{aligned} \quad (36)$$

and the normalization factor K is chosen in such a way that $\mathcal{U}(1) = 1$. Here the sum is over S_i^j and $\bar{S}_i^j$ that are the elements of the following tables:

$$\begin{array}{cccccc} S_1^1 & \dots & S_i^1 & \dots & S_{q_1}^1 & \dots \\ S_1^2 & \dots & S_{q_2}^2 & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ S_1^{c_1} & \dots & \dots & \dots & \dots & \dots \end{array} \quad \begin{array}{cccccc} \bar{S}_1^1 & \dots & \bar{S}_i^1 & \dots & \bar{S}_{q_1}^1 & \dots \\ \bar{S}_1^2 & \dots & \bar{S}_{q_2}^2 & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \bar{S}_1^{c_1} & \dots & \dots & \dots & \dots & \dots \end{array} \quad (37)$$

(The lines contain $q_1, q_2, \dots, q_{c_1}$ elements while the columns contain $c_1, c_2, \dots, c_{q_1}$ elements respectively.) Each of these two tables is obtained via the composition of some permutation inside the lines and some permutation inside the columns of the following table:

$$\begin{array}{cccc} 1 & \dots & 1 & \dots & 1 \\ 2 & \dots & 2 & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \\ c_1 & \dots & \dots & \dots & \dots \end{array} \quad (38)$$

$P(S) = 0(1)$ if the correspondent column permutation is even (odd).

We can rewrite (36) in a compact form

$$\mathcal{U}(U) = \omega_{c_1}(U)\omega_{c_2}(U)\dots\omega_{c_{q_1}}(U), \quad (39)$$

where

$$\omega_M(U) = \det \begin{pmatrix} U_{11} & \dots & U_{1M} \\ \dots & \dots & \dots \\ U_{M1} & \dots & U_{MM} \end{pmatrix} \quad (40)$$

In order to demonstrate the $U(1)$ nature of the derived expression for the Wilson loop (34) we rewrite it as follows:

$$W_q[\mathcal{C}] = \int Dg \exp(i(\mathcal{A}^g, j(\mathcal{C})) + \log(D(q))|\mathcal{C}|), \quad (41)$$

where

$$\mathcal{A}_{xy}^g = -i \log \mathcal{U}(U_{xy}^g) \quad (42)$$

is a complex - valued link field. Below we shall show that its real part can be treated as a lattice Abelian gauge field. (It is worth mentioning that its imaginary part vanishes in a continuum limit. This will be shown in the next section, when we shall consider the continuum limit of the derived expressions.)

The exponent in (41) is invariant under the $U(N - c_1)$ - transformation

$$U_{xy} \rightarrow h^+(\Omega_x) U_{xy} h(\Omega_y) \quad (43)$$

where $\Omega_x \in U(N - c_1)$ is attached to the points of the path $\mathcal{C}$,

$$h(\Omega) = \begin{pmatrix} e^{i\alpha} & 0 & \dots & \dots & \dots \\ 0 & e^{i\alpha} & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & 0 & e^{i\alpha} & 0 \\ \dots & \dots & \dots & 0 & \Omega \end{pmatrix}, \quad (44)$$

and

$$\det \Omega = e^{-ic_1 \alpha} \quad (45)$$

This transformation acts on $\mathcal{A}^g$ as follows:

$$\mathcal{A}_{xy}^g \rightarrow \mathcal{A}_{xy}^{hg} = \mathcal{A}_{xy}^g + r(\alpha_x - \alpha_y), \quad (46)$$

where $r = \sum_i q^i$ is the rank of the representation. This demonstrates that the mentioned $U(N - c_1)$ symmetry induces the $U(1)$ gauge symmetry and real part of $\mathcal{A}^g$ plays the role of the correspondent gauge field.

In analogy with the continuum theory we obtain the lattice variant of the non - Abelian Stokes theorem:

$$W_q[\mathcal{C}] = \int Dg \exp(i(d\mathcal{A}^g, m(\mathcal{C})) + \log(D(q))|\mathcal{C}|), \quad (47)$$

where $m(\mathcal{C})$ is the integer - valued plaquette variable that represents the surface that spans the contour $\mathcal{C}$. $d\mathcal{A}^g$ is the plaquette variable that is defined in an analogy with the Abelian field strength:

$$[d\mathcal{A}]_{xyzw} = \mathcal{A}_{xy} + \mathcal{A}_{yz} - \mathcal{A}_{wz} - \mathcal{A}_{xw}, \quad (48)$$

where we suppose that the directions (xy) and (yz) are positive. The sum over the plaquettes is denoted by

$$(m, d\mathcal{A}) = \sum_{xyzw} m_{xyzw} [d\mathcal{A}]_{xyzw} \quad (49)$$

Let us mention two particular cases, when expression (40) has the most simple form.

For the symmetric ($q_1 = r, q_2 = \dots = q_{N-1} = 0$) representations we have: $\mathcal{U} = U_{11}^r$. Thus the expression for the Wilson loop has the form:

$$W_r[\mathcal{C}] = \int Dg \exp(r(\log U_{11}^g, j(\mathcal{C})) + \log(D(q))|\mathcal{C}|) \quad (50)$$

(The adjoint representation is an example of such a representation. It corresponds to $r = 2$.)

Another simple case corresponds to the absolutely antisymmetric representation ($q_1 = q_2 = \dots = q_r = 1, q_{r+1} = \dots = q_{N-1} = 0$). In this case we have:

$$\mathcal{U} = \det \begin{pmatrix} U_{11} & \dots & U_{1r} \\ \dots & \dots & \dots \\ U_{r1} & \dots & U_{rr} \end{pmatrix} \quad (51)$$

V. THE CONTINUUM LIMIT.

In order to consider consistently the continuum limit of a lattice model, its behavior in a small vicinity of the critical couplings should be investigated. Only near the point of the second order phase transition the lattice correlation length becomes infinite and the lattice spacing can be considered as infinitely small. In the correspondent model the continuum gauge field that is not smooth may arise. It was shown, for example, that such singular configurations of a gauge field as the center vortices may survive a continuum limit [6].

In this paper we consider only smooth continuum gauge fields. The continuum limit of the derived lattice expressions in the case of singular field configurations is out of scope of the present consideration.

Let us consider the expression (41) within the lattice $SU(N)$ gauge model that has the second order phase transition. Let us also consider the behavior of the following correlator:

$$\langle O(U)W_q[\mathcal{C}] \rangle \quad (52)$$

Here $O(U)$ is gauge invariant functional. We assume that $\mathcal{C}$ has no selfintersections. We have:

$$\langle O(U)W_q[\mathcal{C}] \rangle = \langle O(U) \exp(i(\mathcal{A}, j(\mathcal{C})) + \log(D(q))|\mathcal{C}|) \rangle, \quad (53)$$

Here the integration over gauge transformations was absorbed by the average $\langle \dots \rangle$. Let us suppose that the singular gauge fields do not arise in the continuum limit of the considered model. Then we can use the correspondence:

$$U_{xy} \rightarrow (1 + iA_\mu \Delta x), \quad (54)$$

where A is a continuum gauge field and Δx is an infinite small space distance. In accordance with (54) $\langle 0|\mathcal{D}_q(U)|0 \rangle$ corresponds to

$$\langle 0|\mathcal{D}_q(U_{xy})|0 \rangle \rightarrow 1 + i\langle 0|\mathcal{D}_q(A_\mu)|0 \rangle \Delta x \quad (55)$$

Hence it is clear that

$$\mathcal{A}_{xy} \rightarrow \langle 0|\mathcal{D}_q(A_\mu)|0 \rangle \Delta x \quad (56)$$

and imaginary part of $\mathcal{A}$ vanishes in a continuum limit.

Recall that in the vicinity of the second order phase transition $\mathcal{C} = \mathcal{C}_{\text{phys}} \frac{1}{a}$, where the lattice spacing a tends to zero and the physical length of the contour is equal to $\mathcal{C}_{\text{phys}} = \int_{\mathcal{C}} \sqrt{dx_\mu dx_\mu}$. Let us consider such O that it tends to a smooth functional of A in the continuum limit. Therefore we can write the continuum limit of (53) as follows:

$$\begin{aligned} \langle O(U)W_q[\mathcal{C}] \rangle &= \langle O(A) \exp(i \int \langle 0|\mathcal{D}_q(A_\mu)|0 \rangle dx_\mu \\ &\quad + m_q \int_{\mathcal{C}} \sqrt{dx_\mu dx_\mu}) \rangle \end{aligned} \quad (57)$$

where the constant m_q is divergent as

$$m_q \sim \mathbf{M} \log D(q), \quad (58)$$

Here $\mathbf{M}$ is the ultraviolet cut-off (which in the case of the lattice regularization is equal to the inverse lattice spacing).

In gauge theory any Wilson loop always corresponds to the charged particle. The divergence of m_q in (61) should be included into the renormalization scheme for the mass of this particle.

Let us restore the integration over the gauge transformations. The obtained continuum theory is gauge invariant. Hence we can rewrite the last expression as:

$$\begin{aligned} \langle O(U)W_q[\mathcal{C}] \rangle &= \langle O(A) \int Dg \exp(i \int \langle 0|\mathcal{D}_q(A_\mu)|0 \rangle dx_\mu \\ &\quad + m_q \int_{\mathcal{C}} \sqrt{dx_\mu dx_\mu}) \rangle, \end{aligned} \quad (59)$$

where Dg is the measure on the space of all *smooth* gauge transformations, defined in such a way that $\int Dg = 1$.

Due to the arbitrariness of the average $\langle \dots \rangle$ we have:

$$W_q[\mathcal{C}] = \int D\mu_{\mathcal{C}}(g) \exp(i \int \langle 0|\mathcal{D}_q(A_\mu)|0 \rangle dx_\mu), \quad (60)$$

where

$$D\mu_{\mathcal{C},q}(g) = \mathcal{F}(\mathcal{C}) Dg = \exp(m_q \int_{\mathcal{C}} \sqrt{dx_\mu dx_\mu}) Dg \quad (61)$$

This measure differs from the conventional one, which is normalized in such a way that $\int Dg = 1$. Formally $\mathcal{F} \rightarrow \infty$ in the continuum limit. Thus if we substitute Dg into (2) instead of $D\mu_{\mathcal{C},q}(g)$, then the integral vanishes identically in the continuum limit. (This vanishing was pointed out in [7] for some particular choices of A .)

Of course, in any further application of this formula we should remember that the integration is over the smooth gauge transformations and the formula itself is valid only within the model, in which the singular gauge configurations are forbidden.

We use Cartan representation (10) of A^g and obtain

$$\begin{aligned} \langle 0|\mathcal{D}_q(A^g)|0 \rangle &= \langle 0| \sum_{i=1, \dots, N-1} H^i \text{Tr } A^g H^i |0 \rangle \\ &= \text{Tr } A^g \mathcal{H}^q \end{aligned} \quad (62)$$

Thus the representation (60) has the form (2) originally proposed by D.Djakonov and V.Petrov [3].

Using the expressions for the Cartan elements H^i and for the highest weight m^i it is easy to calculate $\mathcal{H}^q$. It's nonzero elements are:

$$\mathcal{H}_{ii}^q = q_i - \frac{1}{N} \sum_{k=1, \dots, N} q_k \quad (i = 1, \dots, N), \quad (63)$$

where we imply that $q_N = 0$.

Hence, we can rewrite (60) as follows:

$$W_q[\mathcal{C}] = \int D\mu_{\mathcal{C}}(g) \exp(i \int \sum_{i=1, \dots, N-1} ([A_\mu^g]_{ii} q_i) dx_\mu) \quad (64)$$

It is worth mentioning that this expression can be obtained directly from (41) (with $\mathcal{U}$ that is given in (40)).

VI. CONCLUSIONS

In this work we have derived the Abelian representation (41) for the $SU(N)$ Wilson loop on the lattice. This representation contains the complex - valued link field. This field is a function of the matrix elements of the $SU(N)$ gauge field. (41) possesses the $U(1)$ gauge symmetry. Real part of the mentioned link field can be considered as a correspondent gauge field. Its imaginary part vanishes in a continuum limit.

The Non - Abelian Stokes theorem on the lattice commonly follows from (41). It can be used for "counting" of the "magnetic" flux that is contained inside the contour correspondent to the Wilson loop. Let us notice that both in the P - vortex and in the monopole pictures of a confinement phenomenon the Wilson loop is approximated by

certain Abelian Wilson loop. The Abelian magnetic flux inside the correspondent contour appears to be responsible for the confinement. Therefore "counting" of the "magnetic" flux performed with the help of the non - Abelian Stokes theorem can be used in related confinement pictures in order to avoid an inexactitude associated with the mentioned approximation.

It is worth mentioning that the continuum limit of the derived expression coincides with the original representation obtained by D. Diakonov and V. Petrov.

Acknowledgments

The author is grateful to A.N.Ivanov for the discussions, which stimulated him to investigate the subject considered in this paper. This work was partly supported by the grants INTAS 00-00111 and CRDF award RP1-2364-MO-02.

-
- [1] G. 't Hooft Nucl. Phys. **B 190**, 455 (1981).
 - [2] M.I. Polikarpov Nucl. Phys. Proc. Suppl. **53**, 134 (1997).
 - [3] D. Diakonov and V. Petrov, Phys. Lett. **B 224**, 131 (1989).
 - [4] D. Diakonov and V. Petrov, hep-th/0008004.
 - [5] K.-I. Kondo and Y. Taira, hep-th/9911242.
 - [6] B.L.G. Bakker, A. I. Veselov and M. A. Zubkov, Phys. Lett. **B 544**, 374 (2002).
 - [7] M. Faber, A. N. Ivanov, N. I. Troitskaya and M. Zach, Phys. Rev. **D 62**, 025019 (2000).
 - [8] M.I. Polikarpov, U. J. Wiese and M. A. Zubkov, Phys. Lett. **B 309**, 133 (1993).
 - [9] M. Creutz, *Quarks, gluons and lattices*, (Cambridge University Press, Cambridge, 1985).